PROCEEDINGS OF THE FOURTH INTERNATIONAL CONGRESS OF ETHNOBOTANY (ICEB 2005)

21-26 August 2005 Istanbul - Turkey

"Ethnobotany: At the Junction of the Continents and the Disciplines"

Hosted and sponsored by Yeditepe University

Edited by Z. Füsun ERTUĞ

(Offprint)



Proceedings of the IVth International Congress of Ethnobotany (ICEB 2005) Yeditepe University Istanbul, 21-26 August 2005

ISBN: 975-807-153-X © 2006 Ege Yayınları

Edited by Z. Füsun Ertuğ

Graphic design Hülya Tokmak Zero Prod. Ltd.

Cover design Ender Yeşildağ

Printed by Mas Matbaacılık A.Ş.

First Edition November 2006, Istanbul

Procedings of the Fourth International Congress of Ethnobotany (ICEB 2005) Yeditepe University, Istanbul 21-26 August 2005, Z. F. Ertuğ ed., Ege Yayınları, Istanbul, 2006.

All rights reserved. No part of this book may be used or reproduced in any manner without written permission of the publisher and the authors.

Publication and Distribution
Zero Prod. Ltd.

Arslan Yatağı Sok. Sedef Palas, 35/2 Cihangir 34433 Istanbul-Turkey
Tel: +90 (212) 244 75 21 - 249 05 20 Fax: +90 (212) 244 32 09
e-mail: info@zerobooksonline.com
www.zerobooksonline.com

Table of Contents

Table of Contents	iii
Organizers of ICEB 2005	xi
A Brief History of the ICEB	xii
Panel and Workshop Descriptions	xiii
Foreword and Acknowledgements	ζvii
PLENARIES	
Montserrat GISPERT CRUELLS A novel approximation to the alimentary culture from an ethnobotanical perspective	1
Vernon H. HEYWOOD Biodiversity, global change and human health	9
Shillean T. PRANCE Some current challenges facing the indigenous peoples of South America	. 21
Nancy J. TURNER Lessons from the grandmothers: Women's roles in traditional botanical knowledge and wisdom in Northwestern North America	. 27
PANEL 1: Ethnobotanical Studies of Wild Plant Foods	
Achille Ephrem ASSOGBADJO, E. DE CALUWÉ, Brice SINSIN, J.T.C. CODJIA, Patrick VAN DAMME Indigenous knowledge of rural people and importance of baobab tree (<i>Adansonia digitata</i> L.) in Benin	. 39
Manuel PARDO-DE-SANTAYANA, Javier TARDÍO, Ana Maria CARVALHO, Juan José LASTRA, Elia SAN-MIGUEL, Emilio BLANCO, and Ramón MORALES Diversity and selection of wild food plants in six regions of Northwestern Iberian Peninsula (Spain and Portugal)	. 49
Nancy J. TURNER From the roots: Indigenous root vegetables of British Columbia, their management and conservation	. 57
Gisella CRUZ GARCÍA Children's knowledge and valuation of wild food plants: The influence of an educational program with tribal and non-tribal children in Western Ghats, India	. 65
Luís S. DIAS and Alexandra S. DIAS Herbs and spices in traditional recipes of Alentejo (Portugal)	. 69
M. en C. Argelia DÍAZ RICO Ethnobotany of nourishing plants cultivated and utilizated by the ethnic group Mixteco in Tepuente, Guerrero State, Mexico	. 73
udith ESPINOSA-MORENO, Dora CENTURION-HIDALGO, Jaime Gabriel CAZARES-CAMERO Edible weeds associated to agricultural plots in Tacotalpa, Tabasco, Mexico	. 77
an C. FITZPATRICK A study of recognition, transmission, and use of wild-food plants in two Wichí communities of the Argentine Chaco	. 79

Peter GIOVANNINI Management, biology and cultural importance of a wild food species in the Tehuacán Cuicatlán Valley, Mexico: The case of <i>Enterolobium cyclocarpum</i>	83
Müberra KOŞAR, Mehmet KOYUNCU, and K.Hüsnü Can BAŞER Folk use of some wild and cultivated <i>Allium</i> species in Turkey	87
Şinasi YILDIRIMLI and Aslı DOĞRU KOCA Uses of some Turkish <i>Asparagus</i> and <i>Tamus</i> species as food	91
Ersin YÜCEL and Gülçin YILMAZ Consumption ways of some <i>Rumex</i> species as food in Turkey	93
PANEL 2: Ethnobotanical Studies on Medicinal and Aromatic Plants	
Paul HERSCH-MARTINEZ Current challenges and scenarios regarding the integration of medicinal plants popular knowledge and formal health systems in Mexico	95
Narayan P. MANANDHAR Native phytotherapy among rural population of Nepal	101
María Rosa MARTINEZ, María Lelia POCHETTINO, Marta CRIVOS, Carolina REMORINI, and Anahí SY Gathering and circulation of medicinal plants in a pluricultural context (Misiones, Argentina)	107
Muhammad Ibrar SHINWARI and Maryum Ibrar SHINWARI Ethnobotany of medicinal and aromatic plants in Pakistan: An Overview	
Paolo Emilio TOMEI, Rita Elisabetta UNCINI MANGANELLI, Serena TRIMARCHI, and Fabiano CAMANGI Ethnopharmacobotany in Italy: State of knowledge and prospect in the future	123
Duygu F. ALPARSLAN and Ertan TUZLACI The folk medicinal plants of the European part of Turkey	129
Kemal Hüsnü Can BAŞER, Gülendam TÜMEN, Hulusi MALYER, and Neşe KIRIMER Plants used for common cold in Turkey	133
Rachid BELHATTAB, Georgios KALANTZAKIS, and Dimitrios BOSKOU Antioxidant activity and total phenolic content of two plants belonging to the Lamiaceae family: Origanum glandulosum Desf. and Marrubium vulgare L.	139
Gülşah ÇOBANOĞLU, Cenk SESAL, Yıldız AYDIN, Müşerref ÖZEREN MORGAN, and Zeki SEVEROĞLU The antimicrobial and the antifungal effects of some lichens with a potential medical and economic use in Turkey	143
Burcu ELÇİ-TARIKAHYA, Sadık ERİK, and Ziver BERKMAN Some anticarcinogenic plants and their usage in the Güdül District (Ankara-Turkey)	147
José Salvador FLORES GUIDO, Rita VERMONT-RICALDE, and Jesús KANTÚN Aromatic plants and their application in traditional medicine at the Mayan communities of the Yucatan Peninsula	151
Paolo Maria GUARRERA and Maria Lucia LEPORATTI Analogies and divergences in the use of medicinal plants in different areas of Central and Southern Italy	155
Maria Lucia LEPORATTI and Kamel GHEDIRA Popular phytotherapy in Italy and Tunisia, a preliminary comparison	159
Emi OKUYAMA, Samir Kumar SADHU, Shunsuke YAMAMOTO, Haruhiro FUJIMOTO, Masami ISHIBASHI, and Erdem YEŞİLADA Prostaglandin-inhibitory and antioxidant components of <i>Cistus laurifolius</i> , a Turkish medicinal plant	

Plants used as analgesic in the folk medicine of Turkey	167
Johanna PUTSCHER and Christian R. VOGL An ethnobotanical survey on herbal medicine in Quito markets	173
Erika RIVERA-ARCE, X. LOZOYA, R. ALVARADO, E. ZÁRATE, J. AGÜERO, M. CHÁVEZ, and M. GATTUSO Ethnobotany and pharmacognosy of the Mexican Plant Drug: <i>Mimosae tenuiflorae</i>	177
G. Maria SALUD PÉREZ, Miguel A.S. ZAVALA, Daniel M. ZAVALA, and Cuauhtemoc G. PÉREZ Anti-diarrhoeal activity of <i>Chysactinia mexicana</i>	
G. Maria SALUD PÉREZ, Miguel A.S. ZAVALA, Cuauhtemoc G. PÉREZ, and Daniel M. ZAVALA Anti-diarrhoeal effect of different extracts of <i>Bidens odorata</i>	185
Chandrakant B. SALUNKHE and N.K. DRAVID Little known medicinal uses of some flowering plants of Maharashtra, India	189
Cristina P. SÁNCHEZ ROJAS, M. Reyes GONZÁLEZ-TEJERO, José M. RAMIRO GUTIÉRREZ, M. CASARES PORCEL, and J. MOLERO MESA Ethnobotany in Sierra de Huelva (South Spain): Medicinal plants	193
Sara SANTOS, Ana I.D. CORREIA, A. Cristina FIGUEIREDO, Luís S. DIAS, and Alexandra S. DIAS The use of herbal remedies in urban and rural areas of the Setúbal Peninsula (Portugal): A study among elders	
Fatih SATIL, Tuncay DİRMENCİ, and Gülendam TÜMEN The trade of wild plants that are named as <i>Thyme</i> (<i>kekik</i>) collected from Kazdağ	201
Muhammad Ibrar SHINWARI and Maryum Ibrar SHINWARI Ethnobotanical study of medicinal and aromatic plants of moist temperate Himalayas in Pakistan	205
Mitra TAGHIZADEH, Mohamad Reza SALEHI ALEA, and Mahshid TAGHIZADEH The efficacy of a herbal mouthwash on the control of gingivitis	209
Zeynep TUNALIER, Neşe KIRIMER, and K. Hüsnü Can BAŞER Demise of a 60-year old Turkish herbal medicine: Lityazol Cemil	213
Gülendam TÜMEN, Hulusi MALYER, K. Hüsnü Can BAŞER, and Serap ÖZ AYDIN Plants used in Anatolia for wound healing	217
PANEL 3: Plant Use of Farmers and Pastoralists	
Patricia C. ANDERSON Non-mechanised processing and storage of cereals, grasses and pulses used for fodder, fuel, food and crafts: Examples from N. Tunisia, Atlas Region, Northwestern Tell	223
Christian Christian BERTSCH, R. VOGL, and Carolina Joana DA SILVA Ethnoveterinary medicine for cattle and horses in the Northern Pantanal Matogrossense, Brazil	233
Jeanne T. GRADE and Patrick VAN DAMME Goat's self-medication against internal parasites in Karamoja, Uganda	241
Natesan PUNNIAMURTHY Ethno Veterinary Medicine (EVM): Use of fresh herbal extracts under field conditions for primary veterinary health care in India	249
Tedje VAN ASSELDONK and Helen BEIJER Herbal folk remedies for animal health in the Netherlands	257
Manuela MANCA and Tania MANCA Uses of plants by shepherds in working process of milk in the village of Bitti (Sardinia, Italy)	265
Jose MARTÍNEZ GONZÁLEZ and Juan Jose LASTRA MENÉNDEZ Ethnobotanical study of the cattle farmers profile in the Aller Valley (Asturias, Spain)	269

Antonio C. PERDOMO-MOLINA The use of local fig tree varieties (<i>Ficus carica</i> L.) for animal feed purposes in the Canary Islands (Spain)	. 273
Ila SHRESTHA and Keshab SHRESTHA Medicinal plants in ethnoveterinary practices in Langtang National Park, Nepal	. 277
Lucia VIEGI, Simonetta BULLITTA, and Giovanna PILUZZA Traditional veterinary practices in some rural areas of Sardinia (Italy)	. 281
Lucia VIEGI, Ignazio CAMARDA, and Giovanni PIRAS Some aspects of ethnoveterinary medicine in Sardinia (Italy)	. 285
PANEL 4: Reproduction and Transmission of Botanical Knowledge and Technology in Basketry and Plaited Artifact	
Belle ASANTE Reviving sustainable plant-based crafts when recent trends favor synthetic fiber usage: Stylistic vicissitudes of Harari baskets in Ethiopia	. 289
Z. Füsun ERTUĞ An overview of the plaited crafts of Turkey (Anatolia and Thrace)	. 297
Dario NOVELLINO Weaving traditions from Island Southeast Asia: Historical context and ethnobotanical knowledge	. 307
Dario NOVELLINO An account of basket weaving and the use of fibre plants in the Mount Aurunci Regional Park (Central Italy)	. 317
Sonia A. VOUGIOUKALOU Weaving knowledge and weaving plants: What will survive the 21st century?	. 327
Ana Maria CARVALHO, Manuel PARDO-DE-SANTAYANA, and Ramón MORALES Traditional knowledge of basketry practices in a Northeastern Region of Portugal	. 335
Giovanni PIRAS Plant-derived utensils employed in traditional agro-pastoral activities in Northwest Sardinia, Italy	. 339
José M. RAMIRO GUTIÉRREZ, M. Ramirez GONZÁLEZ-TEJERO, and Cristina P. SÁNCHEZ-ROJAS Ethnobotany in Huelva province (Spain): Basketry and woodcarving	. 343
Javier TARDÍO, Laura ACEITUNO, and Ramón MORALES The use of plant-based brooms in the province of Madrid (Spain)	. 347
PANEL 5: Biodiversity and Genetic Resources Francisco BASURTO, Virginia EVANGELISTA, Myrna MENDOZA, and Miguel Angel MARTINEZ ALFARO Food fruits of house and a office plantations in the Siegra Negte de Pueble Manies	251
Food fruits of home gardens and coffee plantations in the Sierra Norte de Puebla, Mexico Nina L. ETKIN Wild plant management in rural Hausaland: Local ecological knowledge contributes to the conservation of biodiversity	
Vernon H. HEYWOOD Human use of plant resources - the knowledge base and conservation needs	. 365
Amin U. KHAN Demonstrating the forgotten values of a threatened thorn forest community at an archaeological site: The case for popularizing values in an illustrative manner	
Miguel Ángel SERRATO CRUZ and Juan Saúl BARAJAS PÉREZ *Tagetes filifolia* Lag.: A Mexican genetic resource	. 377

Rosa María FLORES-SERRANO, J. Salvador FLORES, Rosario ITURBE, and Guillermina PEREZ Plant diversity and soil contamination (hydrocarbons and metals): A case study in Ciudad Madero, Tamaulipas, Mexico	383
PANEL 6: Continuity and Change in Food and Medicine: Archaeobotany and the Written Record	
Sabine BECKMANN Root, resin, red and ritual purification - The role of terebinth in Eastern Mediterranean Bronze Age cult	387
Aylen CAPPARELLI, M. Lelia POCHETTINO, Andreoni DIEGO, and Rubén D. ITURRIZA Differences between written and archaeological record: The case of plant micro remains recovered at a Northwestern Argentinean Pipe	397
Patricia A. CLARK Iatrosophia and oral traditions: A case study in Crete's Amari valley	407
Tracey LU The exploitation of Taro in South China	413
Peter J. MATTHEWS Written records of Taro in the Eastern Mediterranean	419
Fragkiska MEGALOUDI Kollyva and funeral bread offerings in Greece: The example of Thasos Island Rolando M.D. NERI-VELA The Badianus Codex and Ophthalmology	
Alexandra LIVARDA and Georgia KOTZAMANI Plant lore in 'Dark Age' Greece: Archaeobotanical evidence from Lefkandi, Euboea, literal sources and traditional knowledge combined	
Emel OYBAK DÖNMEZ Plant use at Early Bronze Age Gre Virike (Şanlıurfa, Turkey)	439
PANEL 7: Capturing Local Perceptions and Priorities of Ethnobotanical Resources Lilian GONZALEZ-CHEVEZ and Paul HERSCH-MARTINEZ Medicinal ethnobotany and meaning construction. A semiotic analysis of plants through representations and practices of some illnesses from the Nahua culture of Guerrero, Mexico	443
Ramón MORALES, Manuel PARDO-DE-SANTAYANA, and Javier TARDÍO The perception of plants in the complete works of Cervantes, particularly "Don Quijote"	
Manuel PARDO-DE-SANTAYANA and Elia SAN-MIGUEL The gender of plants according to popular nomenclature in the North of Spain	461
Miguel Ángel SERRATO-CRUZ Cempoalxochitl: A Mexican flower symbolizing the human	467
Barbara FRUTH and Musuyu MUGANZA Traditional use of wild rainforest plants by the Nkundo, Democratic Republic of Congo (DRC)	471
PANEL 8: Gender Issues in the Ethnobotanical Research	
Erika FRIEDL Old plants and new woman in the Zagros Mountains, Iran	475
Martina Aruna PADMANABHAN Governing the use and conservation of agricultural biodiversity: Institutional and gender analysis of rice cultivation in South India	483
Begum SHAHEEN and Mohammad Adnan SAHIBZADA Role of women in collection, processing and marketing of medicinal plants in Roringar Valley, Swat, Pakistan	491

Nancy J. TURNER "Those Women of Yesteryear": Woman and production of edible seaweed (<i>Porphyra abbottiae</i>) in Coastal British Columbia, Canada	499
PANEL 9: Conservation and Development: Ethnobotanical Discipline at Ethical and Professional Crossroads	
Anil K. GUPTA Ethical issues in accessing people's knowledge and innovations for developing low cost health technologies	507
Sarah-Lan MATHEZ STIEFEL and Madyo D.N. COUTO Linking applied ethnobotany to social learning: A participatory tool for the promotion of indigenous plants use in Matutuíne, Southern Mozambique	517
PANEL 10: Theory and Methodology in the Study of Ethnobotany	
Montse RIGAT, M. Àngels BONET, Sònia GARCIA, Teresa GARNATJE, Joan VALLÈS Ethnobotanical studies in the High River Ter Valley (Pyrenees, Catalonia, Iberian Peninsula)	523
Ignazio CAMARDA Ethno-systematic of Sardinian flora as a scientific system	527
Dora CENTURION-HIDALGO, Jaime Gabriel CAZARES-CAMERO, Judith ESPINOSA-MORENO, and Alberto MAYO-MOSQUEDA Ethnobotanical study of Arecaceae of Tabasco's Sierra of Mexico	531
Melissa CEUTERICK and Patrick VAN DAMME Indigenous use, nomenclature and classification of plants in a Nahuatl-speaking village in the Balsas-Basin, Guerrero, Mexico	533
César del C. LUNA-MORALES Science, traditional knowledge and ethnobotany	537
Maryum Ibrar SHINWARI and Muhammad Ibrar SHINWARI Nomenclatural ambiguity found about herbal crude drug material used in <i>Unani</i> Medicines of Pakistan	541
PANEL 11: Transmission, Contact and Exchange of Plant Resources and Knowledge Between Region Historical and Contemporary Approaches	s:
Amélia FRAZAO-MOREIRA Arabic Gum: From its historical importance in the global markets to its contemporary significance in the local context of Mauritania	545
Stefanie KLAPPA Sago and the settling of Sahul: how present patterns of plant use may illuminate subsistence prehistory	551
Carlos R. RAMIREZ SOSA Quantitative ethnobotany in El Salvador, Central America: A model to study ethobotanical knowledge dynamics	557
Özge SAMANCI Vegetable Patrimony of the Ottoman Culinary Culture	565
BROAD SPECTRUM STUDIES	
Hugo J. DE BOER and Anneleen KOOL Ethnobotany at Uppsala University	571
Ali A. DÖNMEZ and S. Olga BAYRAKTAROĞLU Plants of the <i>İnce Memed</i> : A novel written by Yaşar Kemal	573
Ali A. DÖNMEZ, Emel OYBAK DÖNMEZ, and Serap IŞIK Cultivation of the saffron (<i>Crocus sativus</i> L.) in Türkiye and saffron stories	577

Gizem EMRE BULUT and Ertan TUZLACI An ethnobotanical study in Bozcaada (Çanakkale-Turkey)	581
Fatma GÜNEŞ	
The ethnobotanical importance of some <i>Lathyrus</i> (Fabaceae) species	585
İbrahim İlker ÖZYİĞİT, Nermin GÖZÜKIRMIZI, and Belma D. SEMİZ Agrobacterium-mediated transformation of cotton: Gossypium hirsutum L. "Çukurova 1518" from Turkey	589
Veerambakkam S. RAMACHANDRAN and Jaya VIJAYAN Contribution to the knowledge of non-timber products from tropical forests: A case study from Walayar Range, Palakkad district, Kerala, India	593
Parviz REZVANI MOGHADDAM and Motlagh AHMADZADEH Agronomical studies of black cumin (<i>Nigella sativa</i>) as a new crop in dry and semi dry agricultural system of Iran	597
Daiva SESKAUSKAITE Plants in Lithuanian Folk Songs	601
Serdar Gökhan ŞENOL, Özcan SEÇMEN, and Emin UĞURLU Some ethnobotanical uses in the rural areas of Ödemiş, Tire, Kiraz (İzmir-Turkey)	605
Tedje VAN ASSELDONK and Arend De HAAS Spontaneous foraging behavior of primates in outdoor enclosures	609
Şinasi YILDIRIMLI, Aslı DOĞRU KOCA, and Muhittin DİNÇ The Turkish folk plant names and some uses	613
CATALOGUE OF THE BASKETRY EXHIBITION	617
Dario NOVELLINO and Z. Füsun ERTUĞ General Introduction "Baskets of the World" the Social Significance of Plaited Crafts	619
Dario NOVELLINO An introduction to basketry in Island Southeast Asia	621
The Philippines (Oriental Mindoro)	626
The Philippines (Southern Palawan)	628
The Philippines (Northern-Central Palawan)	631
Borneo-Indonesia (West/Central Kalimantan)	634
Rajinda K. PURI Indonesia (East Kalimantan)	636
Rachel CHUA Eastern Malaysia (Sabah)	638
Roy F. ELLEN Indonesia, the Moluccas (South Central Seram)	640
Sonia VOUGIOUKALOU An introduction to basketry in Polynesia	643
Polynesia, South Pacific (The Cook Islands)	645
Mary BUTCHER An introduction to basketry in Japan	646
Japan (Kyushu Island)	648

Belle ASANTE An introduction to basketry in Africa	649
Africa (Ethiopia)	653
M. El HAOUZI and Gary J. MARTIN Africa (Morocco)	655
Maurice BICHARD An introduction to basketry in the Mediterranean	658
Dario NOVELLINO Central Italy (Maranola)	664
Francesco MINONNE, Concetta MELE, Antonella ALBANO, and Silvano MARCHIORI Salento (Apulia), Southern Italy	668
Ana Maria CARVALHO Northeastern Portugal (Trás-os-Montes)	669
José M. RAMIRO GUTIÉRREZ, M. Reyes GONZÁLEZ-TEJERO, and Cristina P. SÁNCHEZ ROJAS Spain (Huelva Province)	671
Z. Füsun ERTUĞ Turkey	673
Turkey (Aksaray)	677
Turkey (Buldan-Denizli)	678
Turkey (Bodrum-Muğla)	679
Mesut KIRMACI, Zahide ŞAHİN, and Serhat MANAV Turkey (Aydın)	680
Hasan AKAN Turkey (Mardin)	682
Josephine POWELL Turkey (Anatolia)	683
Mary BUTCHER An introduction to basketry in Eastern and Central Europe	684
Daiva ŠEŠKAUSKAITĖ Lithuania (Sargeliai)	688
Index of the Authors and Organizers	691

Diversity and selection of wild food plants in six regions of Northwestern Iberian Peninsula (Spain and Portugal)

PARDO-DE-SANTAYANA, Manuel^{1,2}, Javier TARDÍO³, Ana Maria CARVALHO⁴, Juan José LASTRA⁵, Elia SAN-MIGUEL², Emilio BLANCO² and Ramón MORALES²

- ¹ Centre for Pharmacognosy and Phytotherapy, The School of Pharmacy, University of London, UK. manuel.santayana@pharmacy.ac.uk
- ² Real Jardín Botánico de Madrid. Plaza de Murillo, 2. E-28014 Madrid. Spain; eliasanmiguel@hotmail.com; emiliobc@teleline.es morales@rjb.csic.es
- ³ Instituto Madrileño de Investigación y Desarrollo Rural, Agrario y Alimentario, Apdo. 127, 28800 Alcalá de Henares, Madrid, Spain javier.tardio@madrid.org
- ⁴ Escola Superior Agraria, Campus de Sta. Apolónia, 5301-855 Bragança, Portugal anacarv@ipb.pt.
- ⁵ Universidad de Oviedo. C/ Catedrático Rodrigo Uría s/n E-33071 Oviedo, Spain jlastra@uniovi.es

This paper compares the traditional knowledge and use of wild edible plants in six rural regions of the Northwest of the Iberian Peninsula. Five of them are in Spain: Campoo, Picos de Europa, Piloña, Sanabria and Caurel and the sixth is in Portugal, Parque Natural de Montesinho. Through semi-structured interviews with local informants, data on the use of 97 species were collected. A semi-quantitative approach was used to document the relative importance of each species and to indicate differences in the selection criteria for consuming wild food species in the regions studied. Social, economic and cultural factors need to be taken into account when trying to understand why some wild edible plants have been consumed while others have not. The data indicate that a high percentage of species are used in most regions (17 species are used in 5 or 6 regions). These shared species include many wild fruit plants (e.g. Rubus ulmifolius Schott, Fragaria vesca L.) and the most popular species of each category of use [e.g. vegetables such as Rumex acetosa L., condiments such as Origanum vulgare L., or plants used to prepare herbal teas such as Chamaemelum nobile (L.) All.].

Key words: wild edible plants, quantitative ethnobotany, Spain, Portugal

Despite the primary reliance of agricultural societies on crop plants, the tradition of consuming wild plants has not been completely eliminated. Many people still gather wild edibles as a supplement of their poor diets while in other places, they still enjoy collecting and consuming them for their flavour, more than for their caloric input. Their nutritional role has been reported in many surveys from around the world (Arenas 2003; Bonet and Vallès 2002; Crowe 2001; Ertuğ 2004; Guarrera 2003; Lockett, Carvelt, and Grivetti 2000; Ogle et al. 2001; Ogoye-Ndegwa and Aagaard-Hansen 2003; Pieroni et al. 2005; Tardío, Pascual, and Morales 2005; Turner 1975). These plants have been important supplements to the diet, providing trace elements, vitamins and minerals.

Most studies on wild edible plants focus on the role of these plants within one culture, one ethnic group and little emphasis has been given to the comparison of food plants in various cultures, ethnic groups or communities (Ladio and Lozada 2003; Díaz-Betancourt et al. 1999). This kind of comparative studies are very useful to understand why edible species are either consumed or rejected.

The aim of this study is to compare the diversity of wild edible plants historically gathered for food purposes in six areas of the Northwest of the Iberian Peninsula (including Spain and Portugal) and to analyze the similarities and differences among them. This comparative method provides interesting insights of the selection criteria for




Fig. 1 Location of the six study regions.

food, and it is useful in determining the cultural importance of a particular plant as opposed to others in the same usage category.

Material and methods

Information was obtained through semi-structured interviews with key informants during the two last decades. Informants with a good traditional knowledge of useful wild plants were sought - mostly elderly people who had lived and worked in the region for many years. Open questions were asked about the consumption of wild foods to gain insight into their present-day use, ways of consumption and preparation, the time of collection and the places where each species was gathered.

For the present study, data were grouped into the following categories of edible plants based on folk perceptions: vegetables, plants whose leaves, stems or even unripe fruits or seeds were consumed; wild fruits or seeds consumed when ripe; home-made liqueurs or other alcoholic drinks; herbal teas, used in general as *digestifs*; plants used for seasoning; finally flowers and roots, both eaten for their sweet flavour. Every plant species

mentioned by one informant within one category of use was counted as one use-report (UR) (see Kufer et al. 2005). For the purposes of the present comparison the species with only one UR at all has been rejected.

All studies were conducted in rural areas of the Northwest of the Iberian Peninsula (five in Spain and one in Portugal): *Campoo* in the South of the province of Cantabria (Pardo-de-Santayana 2003); *Picos de Europa*, a region that includes areas of the provinces of Asturias, Cantabria and León (Lastra 2003; authors personal observations); *Piloña* in central-eastern Asturias (San Miguel 2004); *Caurel*, in the South East of the province of Lugo (Blanco 1996); *Sanabria* in the North West of the province of Zamora (Blanco and Diez 2005) and *Montesinho* in the North East of Portugal (Carvalho 2005), close to Zamora province.

Their landscapes include a mosaic of meadows, forests, rivers and high mountain vegetation growing on varied geological materials and soils. Several types of beech and oak forest [Quercus robur L., Q. petraea (Matt.) Liebl. and Q. pyrenaica Willd.], broom scrubland [Cytisus scoparius

(L.) Link, *C. multiflorus* (L´Hér.) Sweet, *Genista florida* L.], and heath [*Erica cinerea* L., *E. vagans* L., *E. australis* L., *E. umbellata* L., *Calluna vulgaris* (L.) Hull] dominate the landscape.

Until few decades the economy of the regions was based on agriculture, cattle breeding and a number of minor activities. Many fields that were used to grow cereals (for bread), pulses and potatoes have now given way to pasture for cows. While the household economy was largely subsistence-based, additional income was derived from the sale of animals, eggs, butter and handicrafts.

All the studied areas are culturally and biologically rich regions, most included in protected areas, such as National Park of Picos de Europa, Natural Park of Sanabria Lake and Natural Park of Montesinho. These areas are located on the border between the Mediterranean and Eurosiberian floristic regions.

Voucher specimens were deposited at the herbariums of the Royal Botanical Garden of Madrid (MA, Real Jardín Botánico), the University of Oviedo (FCO, Universidad de Oviedo) and the School of Agricultural Engenieering of Bragança (BRESA, Escola Superior Agrária).

Results and discussion

Table 1 shows the overall numerical results of each of the ethnobotanical works and some geographical features of the different surveyed areas. It can be seen that the number of species and the number of URs recorded within these regions varies greatly. Taking into account that a similar methodology was followed in all the ethnobotanical works, this variation on the results might be due to differences in sample size (number of localities visited and informants interviewed), geographical characteristics of the surveyed regions (surface and population), but also to the disparity in knowledge and use of wild edible plants among different human groups.

Table 2 includes the correlations among all of these variables. It shows a clear and positive correlation (r=0.84, p<0.05) between the number of wild edible species obtained in the different areas and the number of informants interviewed and also between the latter variable and the number of localities visited. These facts seems to be quite logical and at least until reaching a maximum value, the higher number of informants are interviewed and the higher number localities are visited,

Table 1. Overall numerical results and geographical features of each surveyed areas

Surveyed areas	N. Species	N. Use-report	Informants	Localities	Surface (km ²)	Population
Campoo	57	474	107	42	1012	23000
Caurel	21	114	39	19	100	1200
Picos de Europa	54	571	131	67	1920	19900
Piloña	34	219	94	51	283	8600
Montesinho	50	905	90	28	786	9172
Sanabria	30	131	44	20	2120	15000

Table 2. Correlation matrix among all the variables

	N. Species	N. Use-report	Informants	Localities	Surface	Population
N. Species	1.00					
N. Use-report	0.77	1.00				
Informants	0.84*	0.62	1.00			
Localities	0.56	0.28	0.90*	1.00		
Surface	0.39	0.11	0.20	0.23	1.00	
Population	0.81	0.29	0.65	0.53	0.70	1.00

Marked correlations (*) are significant at p < 0.05

the higher botanical knowledge will be registered. Although not so significant, there is also a high correlation (r=0.81, p=0.05) between the number of plants used and overall population of each region.

However, in our analysis the number of species is not correlated with the surface of the surveyed area (r=0.39), the contrary as could be expected. A larger area will have in general a richer flora that will probably yield a higher number of useful plants. The cited deviation in the correlation coefficient is clearly originated by the study of Sanabria, because this work is only a preliminary survey with a too small ethnobotanical sample for the great surface of the region, as can be seen in Table 1. This could be demonstrated if we calculate the correlations among the same variables but without the study of Sanabria. That way the correlation coefficient between the number of species and the surface area would be much higher (r=0.79, p=0.11).

Therefore, if only rapid ethnobotanical studies can be made, the best way of obtaining the biggest diversity of plants is to select a large and diverse study area and to visit as many scattered localities as possible in the available time. For the same number of people interviewed, it will certainly provide more plants than if a little area is surveyed. In the former case we can only expect to get a preliminary list of the plants used that will includes the most popular plants of each area, with very little redundant data.

This analysis shows that the size of the sample must be selected to best suit the objectives of the project. If all the plants used for a certain purpose want to be documented, the correct number of informants must be interviewed in order not to waste time and work. The number of informants must be selected according to the population and extension of the study area.

Botanical analysis

The use of 97 plant species, belonging to 25 botanical families, has been registered in all these areas. Regarding to the diversity of the species gathered Rosaceae was the most important family, with 19 species, mainly collected for eating their mature fruits or preparing liqueurs. Other important families were Lamiaceae, with 13 species,

used as condiments and digestive infusions and Asteraceae, with six species ingested as green vegetables or in infusions. Five species of Polygonaceae were mainly consumed as vegetables and other five species of Apiaceae were employed in many categories of use.

All the species gathered were autochthonous except for *Mespilus germanica* L., and *Prunus cerasus* L., which after centuries of cultivation, nowadays grow feral in the area. Many species such as *Corylus avellana* L., *Borago officinalis* L., *Laurus nobilis* L., *Castanea sativa* Mill., *Rubus idaeus* L., *Taxus baccata* L., *Ulmus minor* Mill., *Mespilus germanica* L., *Prunus avium* L., *P. insititia* L., *Ribes uva-crispa* L. and *Origanum vulgare* L. are either collected from the wild or cultivated in gardens.

Plants that have been consumed in five or more of the regions studied, can be considered the most relevant or popular species of the whole surveyed area (Table 3). They include many wild fruits such as nuts (chestnut and hazelnuts, both species which are managed or semi domesticated), berries (such as wild strawberry, blackberry or blueberry), cherries and plums. Other species consumed in most of the regions are those gathered for preparing liqueurs (dwarf cherry and blackthorn), the most appreciated vegetables (watercress, sorrel), condiments (bay leaves, oregano), herbal teas (camomile) and fennel that is used for many purposes.

If we consider the number of URs, the list of plants with the highest number of use-reports is similar than the list of plants that are consumed in most regions. Species with the highest number of URs also include the most important species of each category of use. They do not include any species of certain categories such as flowers, sucked for obtaining their sweet nectar, and roots and other underground sweet parts.

Other species have a quite high number of URs but are only popular in one or two regions. This is the case of some vegetables, such as *Montia fontana* L., consumed raw in salads in Sanabria and Montesinho, or *Bryonia dioica* Jacq., very popular in Montesinho for preparing omelettes, soups and stews.

Nearly all of the most common food species of the region are also widely consumed in the whole Spain and the Mediterranean area. An exception is Rumex acetosa L., which although in Piloña and Campoo is the vegetable most cited, it is not so commonly gathered in the rest of Spain (Tardío, Pardo-de-Santayana, and Morales 2006). However, some edible species that are commonly consumed throughout the Iberian Peninsula such as Asparagus officinalis, Scolymus hispanicus L., Silene vulgaris (Moench) Garcke, Taraxacum officinale Weber, Hypochoeris radicata L., Bryonia dioica Jacq. or Portulaca oleracea L., are not gathered in the regions studied or only seldom collected, although they grow in all of them. On the other hand, species such as Mespilus germanica L., Romulea bulbocodium (L.) Sebast. et Mauri, Sideritis hyssopifolia L., Trifolium alpinum L. are mainly consumed in the Northern regions. This shows that the consumption or rejection of some species, especially wild vegetables, can be used as indicators for establishing cultural "homogeneous or related" areas.

Unusual foods

Some of the food species eaten in these Northwestern regions have been scarcely documented before as food plants in the ethnobotanical literature. That is especially true for the plants used as sweets, either roots or flowers. Crocus nudiflorus Sm., whose sweet roots are eaten, and many plants that were sought for their sweet nectar by children and shepherds such as Pedicularis schizocalyx (Lange) Steininger, Fritillaria pyrenaica L. and Lamium purpureum L. are included in this group of plants scarcely documented. Another interesting species is Halimium lasianthum (Lam.) Spach, whose flowering buds or immature fruits were chewed as a snack. However, in many cases other species of these genera are well-known edible species (Tardío, Pardo-de-Santayana, and Morales 2006; PFAF 2005). These plant uses are frequently ignored by researchers and not considered as proper food. The caloric value of nectar is high and, although many flowers should be sucked to obtain high caloric input, these species can have played a role in human nutrition, especially in children's diets.

But there are also unusual species within other food categories, such as plants used for seasoning. *Physospermum cornubiense* (L.) DC. is used in Montesinho to prepare liqueurs and to flavour

different sorts of cakes made with chestnuts; the leaves and flowers of Salvia sclarea L. are employed in the same region for seasoning soups; and the flowers and young buds of Pterospartum tridentatum (L.) Willk. are still used there to prepare a local dish called "arros malandro", a soggy rice. This plant is also used to flavour game or to give a wild flavour to chicken or rabbit.

Other unusual food species is *Nymphaea alba* L., whose immature raw fruits were eaten in Sanabria in the same way as it was documented in the eastern region of Comunidad Valenciana (Pellicer 2001). The edible use of *Tragopogon pratensis* L., *Rumex obtusifolius* L., *Malva tournefortiana* L. as vegetables are scarcely mentioned before as well.

In other cases, the species are well known in other areas as edible plants but the part consumed or the way of consumption is different. *Scandix australis* L., for instance, was known to be eaten as a vegetable in other Spanish regions (Tardío, Pardo-de-Santayana, and Morales 2006), but not to flavour liqueurs, to obtain aniseed liqueur.

Availability of species

As it was stated before, some edible plants that grow in a certain area are not consumed there. In Caurel, a little and isolated area, less than a third of the available species have been consumed, while in Picos de Europa four fifths of the available species have been consumed. Although isolated regions commonly show a higher knowledge transmitted orally, their isolation also leads to a lack of cultural interchange with other regions. At least at European latitudes, popular knowledge of plants is a mixture of oral traditions transmitted among generations and information provided by books or by foreigners to members of the community either of the living generations or of their ancestors. This fact suggests that cultural and not ecological reasons are responsible for the differences in the selection of species.

Interestingly, there are some differences among the categories of uses. The species used to prepare liqueurs or wild fruits, are usually consumed if available, while flowers or vegetables are commonly not consumed or rejected.

In past times, bad communications, lack of money and the hard climate conditions for growing fruit trees made it very difficult to obtain cultivated fresh fruits. Therefore, people needed to gather it from the wild. In fact, only a little species of wild fruits are rejected in some of the areas, they include such as *Amelanchier ovalis*, rare in the region and the bitter and unpleasant

acorns of *Quercus robur* L. and *Q. petraea* (Matt.) Liebl. However, the low demand for wild vegetables in the surveyed areas could be due to the easy availability of cabbage or lettuce in homegardens.

Table 3. Wild food plants utilized in five or more surveyed areas, with the number of informants that mention each food use. SAN: Sanabria; CAU: Caurel; PIL: Piloña; PIC: Picos de Europa; CPO: Campoo; MON: Montesinho

Family/Species (English name)	SAN	CAU	PIL	PIC	СРО	MON	Food use category	Part used, way of consumption
Apiaceae								
Foeniculum vulgare Mill. (fennel)	3	3	1	2		10	Liqueurs	Aerial part or seeds for liqueurs
				1	1		Vegetables	Tender leaves and stems, raw in salads or stewed
						32	Seasoning	Seeds for seasoning soups and stews
				2		23	Herbal teas	Aerial part or seeds for herbal teas or liqueurs
Asteraceae								
Chamaemelum nobile (L.) All. (Roman camomile)	4	6	26	34	17	13	Herbal teas	Inflorescences, as herbal tea
Betulaceae								
Corylus avellana L. (hazelnut)	2	13	10	34	13	3	Fruits	Fruits, eaten raw, dried, added to cakes
Brassicaceae								
Rorippa nasturtium- aquaticum (L.) Hayek (watercress)	6	3	1		11	28	Vegetables	Tender leaves and stems; raw in salads or stewed in soups
Ericaceae								
Vaccinium myrtillus L. (blueberry)	4		6	22	18		Fruits	Fruits, eaten raw and for making jam
	1	5		6			Liqueurs	Fruits, for making liqueurs
Fagaceae								
Castanea sativa Mill. (chestnut)	12	12	31	19	5	43	Fruits	Fruits, eaten raw, dried, roasted or added as a condiment for stews, sometimes ground into flour
Lamiaceae								
Origanum vulgare L. (oregano)	8	6	10	12	20	48	Seasoning	Inflorescences, for seasoning pork (black pudding, "chorizo", marinated pork), stews and roasted meat
			2		1	2	Herbal teas	Inflorescences, as herbal tea
Thymus pulegioides L. (wild thyme)	2	2		6	3	3	Herbal teas	Inflorescences, as herbal tea
				1		2	Seasoning	Inflorescences, for seasoning fish and meat stews and "chorizo"
Lauraceae								
Laurus nobilis L. (bay tree)	1	7	7	27	12	45	Seasoning	Leaves, for seasoning stews

Polygonaceae								
Rumex acetosa L. (sorrel)	5		22	15	35	18	Vegetables	Basal leaves, raw as a snack, in salads or stewed in soups
Rosaceae								
Crataegus monogyna Jacq. (hawthorn)	4		1	20	17	3	Fruits	Fruits, eaten raw
Fragaria vesca L. (wild strawberry)	2	4	5	21	10	30	Fruits	Fruits, eaten raw or desserts
						25	Liqueurs	Fruits, for making liqueurs
Prunus avium L. (cherry tree)	4	7	1	2	3	37	Fruits	Fruits, eaten raw as dessert
				3		2	Liqueurs	Fruits, for making liqueurs
Prunus cerasus L. (sour cherry tree)						11	Fruits	Fruits, for making jams
	2	3	12	4	5	7	Liqueurs	Fruits, for making liqueurs
Prunus spinosa L. (blackthorn)			6	17	16		Fruits	Fruits, eaten raw after stored
	9	8	9	21	14	6	Liqueurs	Fruits, for making liqueurs
Rosa canina L. (wild rose)			4	4	11		Vegetables	Young shoots, raw as a snack
	6			10	18	3	Fruits	Fruits, eaten raw
				1	1		Liqueurs	Fruits, for making liqueurs
Rubus ulmifolius Schott (blackberry)			5	6	14		Vegetables	Young shoots, raw as a snack
	4	9	7	29	16	27	Fruits	Fruits, eaten raw as dessert or for making jams
			2	4	1	6	Liqueurs	Fruits, for liqueurs

Conclusions

As a result of our comparison, it seems that the patterns of wild plant usage depend mainly on socio-cultural than on biological factors such as climate conditions or the richness of the wild edible flora. Social, economic and cultural factors need to be taken into account when trying to understand why some wild edibles are consumed while others are not. Availability of running water, free time to take care of homegardens, better communications and information exchange, direct contact with nature in everyday life, cultural values, fashion and tastes are some of the facts that explain why wild plants are either consumed or rejected.

Some wild species are still gathered, including those plants that have been historically consumed in all the regions with a high frequency of citation. They are the most important species of each category of use (fruits, vegetables, infusions or liqueurs), grow in all the regions and, in many cases, if not easily available from the wild they are even cultivated. They can be considered "key plants" or the core of wild food flora.

Many of the wild edible plants have been considered to be famine food. Rural people from Spain and Portugal frequently think that these plants are old fashioned, unprofitable, or too time-consuming, and prefer to used cultivated plants or buy their food. However, nowadays some species, such as *Sideritis hyssopifolia* L., are being regarded as local specialties, gourmet food or local food that reflect the regional identity and are becoming more popular.

Acknowledgements

We are grateful to all the people who kindly shared their knowledge and time with us.

Literature Cited

- **Arenas, P.** 2003. Etnografía y Alimentación entre los Toba-Ñachilamole#ek y Wichí-Lhuku'tas del Chaco Central (Argentina). Buenos Aires.
- **Blanco, E.** 1996. El Caurel, las plantas y sus habitantes. Fundación Caixa Galicia, La Coruña.
- ———, and J. Diez. 2005. Guía de la Flora de Sanabria, Carballeda y los Valles. Catálogo de Etnoflora. ADISAC-La Voz, Zamora.
- **Bonet, M.A., and J. Vallès.** 2002. Use of non-crop food vascular plants in Montseny biosphere reserve (Catalonia, Iberian Peninsula). International Journal of Food Sciences and Nutrition 53:225-248.
- Carvalho, A.M. 2005 Etnobotánica del Parque Natural de Montesinho. Plantas, tradición y saber popular en un territorio del Nordeste de Portugal. PhD. Thesis. Universidad Autónoma de Madrid.
- **Crowe, A.** 2001. A field guide to the Native edible plants of New Zealand. Godwit Publishing Limited, Birckenhead, Auckland, New Zealand. 3rd Edition.
- Díaz-Betancourt, M., L. Ghermandi, A.H. Ladio, I.R. López-Moreno, E. Raffaele, and E.H. Rapoport. 1999. Weeds as a source for human consumption. A comparison between tropical and temperature Latin America. Revista de Biología Tropical 47:329-338.
- **Ertuğ, F.** 2004. Wild edible plants of the Bodrum Area (Mugla, Turkey). Turkish Journal of Botany 28:161-174.
- **Guarrera**, **P.M.** 2003. Food medicine and minor nurishment in the folk traditions of Central Italy (Marche, Abruzzo and Latium). Fitoterapia 74:515-544.
- **Kufer, J., H. Förther, E. Pöll, and M. Heinrich.** 2005. Ethnobotany of the Ch'orti'. Journal of Pharmacy and Pharmacology 57:1127-1152.
- **Ladio, A.H. and M. Lozada.** 2003. Comparison of wild edible plant diversity and foraging strategies in two aboriginal communities of Northwestern Patagonia. Biodiversity and Conservation 12:937-951.
- **Lastra, J.J.** 2003. Etnobotánica en el Parque Nacional de Picos de Europa. Ministerio de Medio Ambiente. Parques Nacionales, Madrid.

- Lockett, C.T., C.C. Calvert, and L.E. Grivetti. 2000. Energy and micronutrient composition of dietary and medicinal wild plants consumed during drought. Study of rural Fulani, Northeastern Nigeria. International Journal of Food Sciences and Nutrition 51: 195-208.
- Ogle, B.M., N.N.X. Dung, T.T. Do, and L. Hambraeus. 2001. The contribution of wild vegetables to micronutrient intakes among women. An example from the Mekong Delta, Vietnam. Ecology of Food Nutrition 40:159-184.
- Ogoye-Ndegwa, C., and J. Aagaard-Hansen. 2003. Traditional gathering of wild vegetables among the Luo of Western Kenya A nutritional anthropology project. Ecology of Food Nutrition 42:69-89.
- Pardo-de-Santayana, M. 2003 Las plantas en la cultura tradicional de la antigua Merindad de Campoo. PhD Thesis. Universidad Autónoma de Madrid.
- **Pellicer, J.** 2001. Customari botànic. Recerques etnobotàniques a les comarques centrals valencianes. Edicions del Bullent, 2nd Edition, Picanya.
- **PFAF** 2005. Plants for a Future: edible, medicinal and useful plants for a healthier world. Database, available at http://www.pfaf.org.
- Pieroni, A., S. Nebel, R. Franco Santoro, and M. Heinrich. 2005. Food for two seasons: culinary uses of non-cultivated local vegetables and mushrooms in a south Italian village. International Journal of Food Sciences and Nutrition 56:245-272.
- San Miguel, E. 2004 Etnobotánica de Piloña (Asturias). Cultura y saber popular sobre las plantas en un concejo del Centro-Oriente Asturiano. PhD Thesis. Universidad Autónoma de Madrid.
- **Tardío, J., H. Pascual, and R. Morales.** 2005. Wild food plants traditionally used in the province of Madrid. Economic Botany 59:122-136.
- ——, M. Pardo-de-Santayana, and R. Morales. 2006. Ethnobotanical review of wild edible plants in Spain. Botanical Journal of the Linnean Society 152: 27-71.
- **Turner, N.J.** 1975. Food plants of coastal first peoples. UBC Press, Vancouver.